## **SMART START**

A square has an area of 144 in<sup>2</sup>. What is the length, to the nearest inch, of the diagonal?

- 15 inches
  - 16 inches
  - 17 inches
  - 18 inches
- Brianna starts from home and walks 6 km west. She then takes a left and walks 8 km south.
- Using the shortest distance between her home and current location, how far is Brianna from her home in the scenario described?
  - $\sqrt{2}$  km
  - $0.2\sqrt{7}$ km
  - 10 km
  - 14 km

### Example 1 Use the Distributive Property



Use the Distributive Property to factor each polynomial.

a. 
$$27y^2 + 18y$$

Find the GCF of each term.

$$27y^2 = 3 \cdot 3 \cdot 3 \cdot y \cdot y$$
$$18y = 2 \cdot 3 \cdot 3 \cdot y$$

Factor each term.

Circle common factors.

$$GCF = 3 \cdot 3 \cdot y \text{ or } 9y$$

Write each term as the product of the GCF and the remaining factors. Use the Distributive Property to factor out the GCF.

$$27y^{2} + 18y = 9y(3y) + 9y(2)$$
$$= 9y(3y + 2)$$

Rewrite each term using the GCF. Distributive Property

b. 
$$-4a^2b - 8ab^2 + 2ab$$

$$-4a^2b = -1 \cdot 2 \cdot 2 \cdot a \cdot a \cdot b$$

Factor each term.

$$-8ab^2 = -1 \cdot 2 \cdot 2 \cdot 2 \cdot a \cdot b \cdot b$$

Circle common factors.

$$2ab = 2 \cdot a \cdot b$$

 $GCF = 2 \cdot a \cdot b$  or 2ab

$$-4a^{2}b - 8ab^{2} + 2ab = 2ab(-2a) - 2ab(4b) + 2ab(1)$$
$$= 2ab(-2a - 4b + 1)$$

Rewrite each term using the GCF. Distributive Property

### GuidedPractice

**1A.** 
$$15w - 3v$$
 **3(5w - v)**

1B, 
$$7u^2t^2 + 21ut^2 - ut$$
 ut(7ut + 21t - 1)

## Example 2 Factor by Grouping



Factor 4qr + 8r + 3q + 6.

$$4qr + 8r + 3q + 6$$
 Original expression
$$= (4qr + 8r) + (3q + 6)$$
 Group terms with common factors.
$$= 4r(q + 2) + 3(q + 2)$$
 Factor the GCF from each group.

Notice that (q + 2) is common in both groups, so it becomes the GCF.

$$=(4r+3)(q+2)$$

Distributive Property

### GuidedPractice

**2A.** 
$$rn + 5n - r - 5$$
  $(r + 5)(n - 1)$ 

**2A.** 
$$rn + 5n - r - 5$$
  $(r + 5)(n - 1)$  **2B.**  $3np + 15p - 4n - 20$   $(n + 5)(3p - 4)$ 



## **Example 3** Factor by Grouping with Additive Inverses

Factor 2mk - 12m + 42 - 7k.

$$2mk - 12m + 42 - 7k$$

$$= (2mk - 12m) + (42 - 7k)$$

$$=2m(k-6)+7(6-k)$$

$$= 2m(k-6) + 7[(-1)(k-6)]$$

$$=2m(k-6)-7(k-6)$$

$$=(2m-7)(k-6)$$

Group terms with common factors.

Factor the GCF from each group.

$$6 - k = -1(k - 6)$$

Associative Property

Distributive Property

## GuidedPractice

**3A.** 
$$c - 2cd + 8d - 4$$

**3B.** 
$$3p - 2p^2 - 18p + 27$$

### Example 4 Solve Equations

Solve each equation. Check your solutions.

a. 
$$(2d + 6)(3d - 15) = 0$$

$$(2d + 6)(3d - 15) = 0$$

Original equation

$$2d + 6 = 0$$
 or  $3d - 15 = 0$ 

Zero Product Property

$$2d = -6$$

$$3d = 15$$
 Solve each equation.

$$d = -3$$

$$d = 5$$

Divide.

The roots are -3 and 5.

CHECK Substitute -3 and 5 for d in the original equation.

$$(2d + 6)(3d - 15) = 0$$

$$(2d + 6)(3d - 15) = 0$$

$$[2(-3) + 6][3(-3) - 15] \stackrel{?}{=} 0$$
  $[2(5) + 6][3(5) - 15] \stackrel{?}{=} 0$ 

$$2(5) + 6][3(5) - 15] \stackrel{?}{=} ($$

$$(-6+6)(-9-15)\stackrel{?}{=}0$$
  $(10+6)(15-15)\stackrel{?}{=}0$ 

$$+ 6)(15 - 15) = 0$$

$$(0)(-24)\stackrel{?}{=} 0$$

$$16(0) \stackrel{\circ}{=} 0$$

$$0 = 0$$

$$0 = 0$$

b. 
$$c^2 = 3c$$

$$c^2 = 3c$$

Original equation

$$c^2 - 3c = 0$$

Subtract 3c from each side to get 0 on one side of the equation.

$$c(c - 3) = 0$$

Factor by using the GCF to get the form ab=0.

$$c = 0$$
 or  $c - 3 = 0$ 

Zero Product Property

$$c = 3$$

Solve each equation.

The roots are 0 and 3.

Check by substituting 0 and 3 for a.


#### GuidedPractice

**4A.** 
$$3n(n+2) = 0$$

**4B.** 
$$8b^2 - 40b = 0$$

**4C.** 
$$x^2 = -10x$$

# 8-6 page 503



### Example 1 b and c are Positive

Factor  $x^2 + 9x + 20$ .

In this trinomial, b = 9 and c = 20. Since c is positive and b is positive, you need to find two positive factors with a sum of 9 and a product of 20. Make an organized list of the factors of 20, and look for the pair of factors with a sum of 9.

Factors of 20	Sum of Factors
1, 20	21
2, 10	12
4, 5	9

 $x^2 + 9x + 20 = (x + m)(x + p)$ = (x + 4)(x + 5) The correct factors are 4 and 5.

Write the pattern. m = 4 and p = 5

**CHECK** You can check this result by multiplying the two factors. The product should be equal to the original expression.

$$(x + 4)(x + 5) = x^2 + 5x + 4x + 20$$
 FOIL Method  
=  $x^2 + 9x + 20$  Simplify.

### ▶ GuidedPractice

**1A.** 
$$d^2 + 11d + 24$$

**1B.** 
$$9 + 10t + t^2$$

## Example 2 b is Negative and c is Positive

Factor  $x^2 - 8x + 12$ . Confirm your answer using a graphing calculator.

In this trinomial, b = -8 and c = 12. Since c is positive and b is negative, you need to find two negative factors with a sum of -8 and a product of 12.

Factors	Sum of
of 12	Factors
-1, -12	-13
-2, -6	-8
-3, -4	-7

The correct factors are -2 and -6.

$$x^2 - 8x + 12 = (x + m)(x + p)$$
 Write the pattern.  
=  $(x - 2)(x - 6)$   $m = -2$  and  $p = -6$ 

**CHECK** Graph  $y = x^2 - 8x + 12$  and y = (x - 2)(x - 6) on the same screen. Since only one graph appears, the two graphs must coincide. Therefore, the trinomial has been factored correctly.  $\checkmark$ 



[-10, 10] scl: 1 by [-10, 10] scl: 1

GuidedPractice

**2A.** 
$$21 - 22m + m^2$$

**2B.** 
$$w^2 - 11w + 28$$

## Example 3 c is Negative



Factor each polynomial. Confirm your answers using a graphing calculator.

a. 
$$x^2 + 2x - 15$$

In this trinomial, b = 2 and c = -15. Since c is negative, the factors m and p have opposite signs. So either m or p is negative, but not both. Since b is positive, the factor with the greater absolute value is also positive.

List the factors of -15, where one factor of each pair is negative. Look for the pair of factors with a sum of 2.

Simplify.

Factors of -15	Sum of Factors	
-1, 15	14	
<b>-3</b> , 5	2	The correct factors are -3 and 5.
$x^2 + 2x - 15$	= (x + m)(x + p)	Write the pattern.
	=(x-3)(x+5)	m = -3 and $p = 5$
CHECK $(x-3)$	$3)(x+5) = x^2 + 5x - 3x - 15$	FOIL Method

 $= x^2 + 2x - 15$ 

In this trinomial, b = -7 and c = -18. Either m or p is negative, but not both. Since b is negative, the factor with the greater absolute value is also negative.

List the factors of -18, where one factor of each pair is negative. Look for the pair of factors with a sum of -7.

Factors	Sum of
of -18	Factors
1, -18	-17
<b>2</b> , <b>−9</b>	-7
3, -6	-3


$$x^{2} - 7x - 18 = (x + m)(x + p)$$
$$= (x + 2)(x - 9)$$

The correct factors are 2 and -9.

Write the pattern.

$$m=2$$
 and  $p=-9$ 


**CHECK** Graph  $y = x^2 - 7x - 18$  and y = (x + 2)(x - 9) on the same screen.



# **Guided**Practice


**3A.** 
$$y^2 + 13y - 48$$

**3B.** 
$$r^2 - 2r - 24$$



### **Example 4** Solve an Equation by Factoring





The roots are 3 and -9.

**CHECK** Substitute 3 and -9 for x in the original equation.

$$x^{2} + 6x = 27$$
  
 $(3)^{2} + 6(3) \stackrel{?}{=} 27$   
 $9 + 18 \stackrel{?}{=} 27$   
 $27 = 27 \checkmark$ 
 $x^{2} + 6x = 27$   
 $(-9)^{2} + 6(-9) \stackrel{?}{=} 27$   
 $81 - 54 \stackrel{?}{=} 27$   
 $27 = 27 \checkmark$ 

#### GuidedPractice

Solve each equation. Check your solutions.

**4A.** 
$$z^2 - 3z = 70$$
 **4B.**  $x^2 + 3x - 18 = 0$